



## Diaper Rash Baby Care (O/W)

### Lactokine™ Fluid PF

FORMULA No. 311.02.0950

Phase	Trade Name	INCI Name	w/w %	Supplier
A	Arlacel 165V	Glyceryl Stearate, PEG-100 Stearate	2.00	Croda
	Lanette O	Cetearyl Alcohol	4.00	BASF
	Rice Bran Wax 2811	Oryza Sativa (Rice) Bran Wax	0.40	Kahl
	Liponate DPC-6	Dipentaerythrityl Hexacaprylate/ Hexacaprate	1.50	Vantage
	Isopropyl Palmitate	Isopropyl Palmitate	6.00	BASF
	Phytowax Olive 18 L57	Hydrogenated Olive Oil Stearyl Esters	5.00	Seppic
	Lipex Shea Soft	Soybean Glycerides, Butyrospermum Parkii Butter Unsaponifiables	12.00	AAK
	Dow Corning 1501	Cyclopentasiloxane, Dimethiconol	9.00	Dow Corning
B	Water	Water	ad 100	
	Inutec SP-1	Inulin Lauryl Carbamate	0.80	Beneo
	Sisterna L-70C	Water, Sucrose Laurate, Alcohol	0.20	Sisterna
	Sisterna SP-70C	Sucrose Stearate	0.10	Sisterna
	Glycerin	Glycerin	3.00	
	Veegum	Magnesium Aluminum Silicate	0.50	Vanderbilt
	Vanzan NF	Xanthan Gum	0.15	Vanderbilt
C	<b>Lactokine™ Fluid PF</b>	Milk Protein	<b>3.00</b>	<b>CLR</b>
	Euxyl PE 9010*	Phenoxyethanol, Ethylhexylglycerin	0.80	Schülke
	Tapioca Pure	Tapioca Starch	5.00	Akzo Nobel

#### Manufacture:

Mix A and B. Stir B until completely soaked. Heat up both to 90°C separately. Add A to B and homogenize for 3 minutes. Cool down while stirring and add C in the given order below 30°C.



The recommendations and formulations given are based on our knowledge and experience in the field of technical application. They are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

\*) This formula has been manufactured and stability tested using a special preservative, but has not been subjected to microbiological challenge tests.