



Age Defense Ampoule Treatment

MPC™ - Milk Peptide Complex

FORMULA No. 355.06.0767

| Phase | Trade Name | INCI Name | w/w % | Supplier |
|-------|----------------------------------|------------------------------------|------------|----------------|
| A | Glucam E-20 | Methyl Gluceth-20 | 2.0 | Lubrizol |
| | Tylose H 4000 G4 PHA | Hydroxyethylcellulose | 1.0 | SE Tylose GmbH |
| B | Water | Water | ad 100 | |
| C | Euxyl PE 9010* | Phenoxyethanol, Ethylhexylglycerin | 0.8 | Schülke |
| D | Water | Water | 9.5 | |
| | MPC™-Milk Peptide Complex | Whey Protein | 0.5 | CLR |
| | NaOH (10%) | Water, Sodium Hydroxide | q.s. | |

Manufacture

Heat A to 60°C. Add B and homogenize for 3 minutes. Cool down while stirring and add C at room temperature. Predisperse phase D while adjusting pH value to 6-7. Add D while stirring and mix until homogenous.



The recommendations and formulations given are based on our knowledge and experience in the field of technical application. They are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

*) This formula has been manufactured and stability tested using a special preservative, but has not been subjected to microbiological challenge tests.