



Clear Gel

Elastin CLR, Collagen CLR

FORMULA No. 150.16.0749

Phase	Trade Name	INCI Name	w/w %	Supplier
A	Water	Water	ad 100	
	Pemulen TR-1	Acrylates/C 10-30 Alkyl Acrylate Crosspolymer	0.6	Lubrizol
	Glycerin	Glycerin	3.5	
	Euxyl PE 9010*	Phenoxyethanol, Ethylhexylglycerin	0.8	Schülke
B	NaOH (10%)	Water, Sodium Hydroxide	q.s.	
C	Collagen CLR	Soluble Collagen	5.0	CLR
	Elastin CLR	Hydrolyzed Elastin	5.0	CLR

Manufacture

Mix A and stir until completely dispersed. Adjust pH value to 5.5-6 with B. Add C in the given order and stir until uniform.



The recommendations and formulations given are based on our knowledge and experience in the field of technical application. They are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

*) This formula has been manufactured and stability tested using a special preservative, but has not been subjected to microbiological challenge tests.