



Endless Summer Glow BB Cream (O/W)

Modukine™, Belides™ NP, Vitamin F forte, SyriCalm™CLR

FORMULA No. 380.001.01

Phase	Trade Name	INCI Name	w/w %	Supplier
A	Montanov 68	Cetearyl Alcohol, Cetearyl Glucoside	2.0	Seppic
	Montanov 202	Arachidyl Alcohol, Behenyl Alcohol, Arachidyl Glucoside	2.0	Seppic
	Tegosoft AC	Isoamyl Cocoate	5.0	Evonik
	Cutina FS 45	Stearic Acid, Palmitic Acid	1.0	BASF
	Uvinul A Plus B	Ethylhexyl Methoxycinnamate, Diethylamino Hydroxybenzoyl Hexyl Benzoate	10.0	BASF
	Tinosorb S	Bis-Ethylhexyloxyphenol Methoxyphenyl Triazine	1.0	BASF
	Vitamin F forte	Linoleic Acid, Linolenic Acid	1.0	CLR
	Dermofeel Toco 70 Non GMO	Tocopherol, Helianthus Annuus (Sunflower) Seed Oil	0.05	dr. straetmans
B	Water	Water	ad 100	
	Keltrol CG-RD	Xanthan Gum	0.2	CP Kelco
	Mirage Bright Silver	Titanium Dioxide, Calcium Sodium Borosilicate	0.2	Eckart
	Mirage Sparkling Luxury Gold	Titanium Dioxide, Iron Oxides, Calcium Sodium Borosilicate, Silica	3.8	Eckart
	RonaFlair LDP White	Sodium Potassium Aluminum Silicate, CI 77891	2.0	Merck
	SynCrystal Copper	Synthetic Fluorophlogopite, Iron Oxides	1.5	Eckart
C	Modukine™	Lactose, Milk Protein	0.5	CLR
	Trisodium Citrate Dihydrate	Sodium Citrate	0.12	
	Water	Water	9.38	
D	Belides™ NP	Bellis Perennis (Daisy) Flower Extract	2.0	CLR
	SyriCalm™ CLR	Water, Phragmites Kharka Extract, Poria Cocos Extract	3.0	CLR
	Euxyl PE 9010*	Phenoxyethanol, Ethylhexylglycerin	0.8	Schülke
	NaOH (10%)	Water, Sodium Hydroxide	0.15	



The recommendations and formulations given are based on our knowledge and experience in the field of technical application. They are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

*) This formula has been manufactured and stability tested using a special preservative, but has not been subjected to microbiological challenge tests.



Manufacture

Predisperse B and stir until uniform. Heat up A and B separately to 85°C. Add B to A while stirring and homogenize for 3 minutes. Cool down while stirring and add predispersed phase C below 30°C. Add D in the given order and stir until homogenous.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application. They are, to the best of our belief, correct, but are offered without obligation. Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

*) This formula has been manufactured and stability tested using a special preservative, but has not been subjected to microbiological challenge tests.