



Baked Eyeshadow

** (developed & produced by Eckart; www.eckart.net)

MPC™-Milk Peptide Complex, Glycoderm™ (P)

FORMULA No. 205.01.0958

Phase	Trade Name	INCI Name	w/w %	Supplier
A	Optigel CL-PC	Magnesium Aluminum Silicate	3.00	BYK
	Deionized Water	Water	q.s.	
B	Myritol 318	Caprylic/Capric Triglyceride	3.00	BASF
	Ceraphyl 847	Octyldodecyl Stearoyl Stearate	3.00	Ashland
	Xiameter PMX-200 Silicone Fluid 5cs	Dimethicone	3.00	Dow
	Lipo GMS-450 VEG	Glyceryl Stearate	2.00	Vantage
C	Talc	Talc	8.50	VWR
	Orgasol 2002 D NAT COS	Nylon-12	8.00	Arkema Orgasol
	SYNAFIL S 1050	Synthetic Fluorphlogopite	10.00	Eckart
	Citric Acid (10%)	Water, Citric Acid	1.50	
D	MPC™-Milk Peptide Complex	Whey Protein	0.50	CLR
	Glycoderm™ (P)	Water, Honey, Phospholipids, Sphingolipids, Hyaluronic Acid	2.00	CLR
	MIRAGE Glamour Silver	Calcium Sodium Borosilicate, Titanium Dioxide, Tin Oxide	5.00	Eckart
	MIRAGE Glamour Gold	Calcium Sodium Borosilicate, Titanium Dioxide, Tin Oxide	5.00	Eckart
	SYNCRYSTAL Sapphire	Synthetic Fluorphlogopite, Titanium Dioxide, Iron Oxides, Ferric Ferrocyanide, Tin Oxide	12.00	Eckart
	SYNCRYSTAL Jade	Synthetic Fluorphlogopite, Titanium Dioxide, Iron Oxides, Ferric Ferrocyanide, Tin Oxide	12.00	Eckart

Manufacture

Prepare a 3% suspension of Optigel CL-PC and water under high shear. Heat up phase A and B separately to 75°C. Add B to A under high shear. Add C step by step to AB under high shear. Adjust pH-value to 6-7 with Citric Acid (10%). Add D to ABC under agitation. Pour the suspension to an appropriate mould and let it dry.