



Pore Normalizing Primer Base OW
I-71.385.01A

<i>Phase</i>	<i>Material Name</i>	<i>US INCI</i>	<i>Supplier</i>	<i>% Material</i>
A	UniqSens SFE System	Pectin Xanthan Gum Carrageenan	CP Kelco	2.10
A	Eutanol G	Octyldodecanol	BASF	3.00
A	Perfume Luxury Care	Fragrance	Voegele	0.25
B	Deionized Water	Water	N.A.	81.67
B	MultiMoist CLR™	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	3.00
B	ProRenew Complex CLR™	Lactococcus Ferment Lysate	CLR	3.00
B	CutiFine CLR™	Water Vaccaria Segetalis (Cowherb) Extract Adansonia Digitata (Baobab) Pulp Extract	CLR	3.00
B	Euxyl K 712	Sodium Benzoate Potassium Sorbate Water	Ashland	0.80
C	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.18
D	RonaFlair Flawless	Silica Titanium Dioxide Iron Oxides	Merck	3.00
				100.00

Operating Instructions

Mix A and B separately. Add B to A while stirring and homogenize until uniform. Adjust pH value to 5-5.4 with C and add D as desired.

Directions for use:

Take a pea-sized amount and gently massage into the skin.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com