



SOS Anti - Irritation Gel
II-17.469.01

Phase	Material Name	US INCI	Supplier	% Material
A	Deionized Water	Water	N.A.	80.59
A	Phenethylalcohol nat.	Phenylethyl Alcohol	Evonik	0.60
A	Carbopol Ultrez 21	Acrylates/C10-30 Alkyl Acrylate Crosspolymer	Lubrizol	0.50
A	Hispagel 200 Ns	Glycerin Glyceryl Polyacrylate	BASF	5.00
B	SyriCalm™ CLR (PC)	Water Phragmites Communis Extract Poria Cocos Extract	CLR	3.00
C	Deionized Water	Water	N.A.	9.20
C	Modukine™	Lactose Milk Protein	CLR	0.50
D	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.61
				100.00

Operating Instructions

Mix A and stir slowly until completely dispersed. Add B and premixed C under medium stirring. Adjust pH to 6.5 with D.

Directions for use:

Take a pea-sized amount of the gel and gently massage into the skin.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com