



Anti Redness Roll-On  
II-32.311.01\_2

<b>Phase</b>	<b>Material Name</b>	<b>US INCI</b>	<b>Supplier</b>	<b>% Material</b>
A	Deionized Water	Water	N.A.	88.85
A	Zemea	Propanediol	DuPont Tate & Lyle	2.00
A	Euxyl PE9010	Phenoxyethanol Ethylhexylglycerin	Ashland	0.80
A	Pemulen TR-2	Acrylates/C10-30 Alkyl Acrylate Crosspolymer	Lubrizol	0.20
B	<b>Hygroplex™ HHG</b>	Hexylene Glycol Fructose Glucose Sucrose Urea Dextrin Alanine Glutamic Acid Aspartic Acid Hexyl Nicotinate	CLR	3.00
B	<b>Lactokine™ Fluid PF</b>	Milk Protein	CLR	5.00
C	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.15
				<b>100.00</b>

#### **Operating Instructions**

Mix A and stir until completely dispersed. Add B and stir until homogeneous. Adjust pH value to 6.5 with C.

Directions for use:

Apply the roll-on the reddened and itching skin.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - [www.clr-berlin.com](http://www.clr-berlin.com)