



Stimulating Hair Growth Treatment-Leave-On
II-33.421.03A

<i>Phase</i>	<i>Material Name</i>	<i>US INCI</i>	<i>Supplier</i>	<i>% Material</i>
A	Deionized Water	Water	N.A.	60.58
A	Hispagel 200 Ns	Glycerin Glyceryl Polyacrylate	BASF	25.00
B	MultiMoist CLR™	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	1.00
B	ProRenew Complex CLR™	Lactococcus Ferment Lysate	CLR	3.00
B	Pentylene Glycol	Pentylene Glycol	Cosnaderm	5.00
B	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.27
C	Soluvit Richter NP	Water Alcohol PEG-35 Castor Oil Polysorbate 20 Aesculus Hippocastanum (Horse Chestnut) Seed Extract Retinyl Palmitate Tocopherol Inositol Calcium Pantothenate Linoleic Acid Biotin	CLR	5.00
D	Perfume Grape PMF WS	Fragrance	Voegele	0.15
				100.00

Operating Instructions

Mix A and stir until homogeneous. Add B in this order and adjust pH value to 6-6.5. Add C as desired and stir until uniform.

Directions for use:

Take a pea-sized amount and gently massage into the scalp after hair wash and leave-on.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com