



Pore Refining Face Emulsion WO
III-10.136.05A

Phase	Material Name	US INCI	Supplier	% Material
A	Deionized Water	Water	N.A.	71.77
A	Sepinov EMT 10	Hydroxyethyl Acrylate/Sodium Acryloyldimethyl Taurate Copolymer	SEPPIC	1.25
A	Sepimax ZEN	Polyacrylate Crosspolymer-6	SEPPIC	0.45
B	Glycerol 85%, vegetable Ph. Eur.	Glycerin	Gustav Heess	5.00
B	Nipaguard SCE	Sorbitan Caprylate Propanediol Benzoic Acid	Clariant	0.80
B	NaOH (10%)	Water Sodium Hydroxide	N.A.	0.20
B	ProRenew Complex CLR™	Lactococcus Ferment Lysate	CLR	3.00
B	Belides™ NP	Bellis Perennis (Daisy) Flower Extract	CLR	2.00
B	CutiFine CLR™	Water Gypsophila Vaccaria Flower/Leaf/Stem Extract Adansonia Digitata (Baobab) Pulp Extract	CLR	3.00
C	Easynov	Octyldodecanol Octyldodecyl Xyloside PEG-30 Dipolyhydroxystearate	SEPPIC	3.00
C	Eutanol G	Octyldodecanol	BASF	3.00
C	Lanol 99	Isononyl Isononanoate	SEPPIC	6.00
C	Aromatic White Peach Extract Blend N33008	Available on request	Carrubba	0.53
				100.00

Operating Instructions

Mix A and stir until homogeneous. Add B in this order and adjust pH value to 5.5-6, if necessary. Add premixed phase C to AB while stirring. Stir until uniform (800rpm with propeller stirrer).

Directions for use:

Take a pea-sized amount of the lotion and gently massage into the skin.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com