



Quick Breaking Lotion WO  
III-4.385.01

<i>Phase</i>	<i>Material Name</i>	<i>US INCI</i>	<i>Supplier</i>	<i>% Material</i>
A	Deionized Water	Water	N.A.	74.69
A	Euxyl PE9010	Phenoxyethanol Ethylhexylglycerin	Ashland	0.80
A	Sodium Chloride	Sodium Chloride	N.A.	0.50
A	<b>MultiMoist CLR™</b>	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	3.00
A	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.11
B	Myritol 318 MB	Caprylic/Capric Triglyceride	BASF	20.00
B	Abil EM 180	Cetyl PEG/PPG-10/1 Dimethicone	Evonik	0.90
				<b>100.00</b>

***Operating Instructions***

Mix A and B separately. Slowly add A to B and while stirring fast.

Directions for use:

Use on cleansed skin. Gently massage into the skin.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - [www.clr-berlin.com](http://www.clr-berlin.com)