



Natural Primer Base WO
III-9.138.02A

Phase	Material Name	US INCI	Supplier	% Material
A	Deionized Water	Water	N.A.	66.53
A	Glycerol 85%, vegetable Ph. Eur.	Glycerin	Gustav Heess	5.00
A	Euxyl K 900	Benzyl Alcohol Ethylhexylglycerin Tocopherol	Ashland	0.60
A	CutiGuard CLR™	Betaine Sucrose Hydrolyzed Rhodophyceae Extract Water	CLR	3.00
A	ProRenew Complex CLR™	Lactococcus Ferment Lysate	CLR	3.00
A	MultiMoist CLR™	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	3.00
B	Spherilex 10 PC	Hydrated Silica	Evonik	1.00
B	Sepilife NUDE	Sodium Acrylate/Sodium Acryloyldimethyl Taurate Copolymer C15-19 Alkane Polyglyceryl-6 Laurate Polyglycerin-6	SEPPIC	3.60
C	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.12
D	Easynov	Octyldodecanol Octyldodecyl Xyloside PEG-30 Dipolyhydroxystearate	SEPPIC	2.80
D	Lanol 99	Isononyl Isononanoate	SEPPIC	6.00
E	Mearlmica DD	Mica	BASF	5.00
F	Perfume Nat. Creme de Luxe	Fragrance	Frey+Lau	0.35
				100.00

Operating Instructions

Mix A and stir until homogeneous. Adjust pH value to 6 with C and add D. Stir until uniform (800rpm with propeller stirrer). Add E and F as desired.

Directions for use:

Take a pea-sized amount and gently massage into skin in the morning.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com