



Derma IQ Shiny Hydra Intense Peel-Off Mask
VI-4.421.02A

<i>Phase</i>	<i>Material Name</i>	<i>US INCI</i>	<i>Supplier</i>	<i>% Material</i>
A	Deionized Water	Water	N.A.	62.32
A	Propylene Glycol	Propylene Glycol	BASF	5.00
B	Keltrol CG-BT	Xanthan Gum	CP Kelco	1.00
B	Gohsenol EG-40C	Polyvinyl Alcohol	Mitsubishi Chemicals	10.00
C	Ethanol (96%)	Alcohol	Berkel AHK	15.00
D	ProRenew Complex CLR™	Lactococcus Ferment Lysate	CLR	3.00
D	MultiMoist CLR™	Fructooligosaccharides Beta Vulgaris (Beet) Root Extract Water	CLR	3.00
E	NaOH (32%)	Water Sodium Hydroxide	N.A.	0.18
F	Perfume Yuzu Rage	Fragrance	IFF	0.20
G	Sachet MV Gold S2317	Synthetic Fluorphlogopite Titanium Dioxide Manganese Violet Triethoxycaprylylsilane Tin Oxide	Sandream Impact	0.30
				100.00

Operating Instructions

Mix A and heat up to 85°C. Add B while stirring and heating; keep temperature for 30-60 minutes. Stir until completely dispersed and uniform. Cool down while stirring. Add C and D in the given order below 30°C. Mix until homogeneous. Adjust pH value to 6-6.5 with E. Add F and G as desired.

Directions for use:

Apply to dry skin and leave for about 10-15 minutes (until the mask dries and lifts at the edges). Peel-off mask from one side.

The recommendations and formulations given are based on our knowledge and experience in the field of technical application.

They are, to the best of our belief, correct, but are offered without obligation.

Those who use our recommendations and formulations as well as those who process CLR Active Agents are themselves responsible for the adherence to prevailing statutory regulations and the observance of patent rights as well as other protective rights for other companies.

This formula has been manufactured and stability-tested using a special preservative, but has not been subjected to microbiological challenge tests.

CLR - Chemisches Laboratorium Dr. Kurt Richter GmbH - www.clr-berlin.com